

## DRUG FACTS

**Active ingredient(s)**

**Purpose**

### Use(s)

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages.

### Warnings

**Allergy alert** Do not use if you are allergic to famotidine or other acid reducers.

#### Do not use

#### Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

#### Ask a doctor before use if you have

#### Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating, or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

#### Ask a doctor or pharmacist before use if you are

#### When using this product

#### Stop use and ask a doctor if

#### Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

#### If pregnant or breast-feeding

#### Keep out of reach of children

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

### Directions

- adults and children 12 years and over:
- to **relieve** symptoms, swallow 1 tablet with a glass of water. Do not chew.
- to **prevent** symptoms, swallow 1 tablet with a glass of water at any time from **15 to 60 minutes before** eating food or drinking beverages that cause heartburn
- do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

### Other information

- read the directions and warnings before use
- **TAMPER-EVIDENT:** Do not use this product if carton has been opened, if blister backing appears to have been disturbed, or if individual blister units are broken or torn.
- keep the carton. It contains important information.
- store at 20° to 25°C (68° to 77°F)
- protect from moisture

#### Drug Facts

### Inactive ingredients

colloidal silicon dioxide, FD & C Red #40 aluminum lake, FD & C Yellow #6 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch, sodium starch glycolate, talc, titanium dioxide